

Interindividual and topographical variation of L:M cone ratios in monkey retinas

Samir S. Deeb

Departments of Medicine and Genetics, University of Washington, Seattle, Washington 98195

Lisa C. Diller

Department of Biological Structure, University of Washington, Seattle, Washington 98195

David R. Williams

Center for Visual Sciences, University of Rochester, Rochester, New York 14627

Dennis M. Dacey

Department of Biological Structure, University of Washington, Seattle, Washington 98195

Received July 21, 1999; revised manuscript received November 1, 1999; accepted November 3, 1999

We analyzed the ratio of L:M cone photopigment mRNA in the retinas of Old World monkeys, using the method of rapid polymerase chain reaction–single-strand conformation polymorphism. The L:M cone pigment mRNA ratio in whole retina ranged from 0.6 to 7.0, with a mean of ~ 1.6 (standard deviation, ± 0.56 ; $n = 26$). There was no change in this ratio with eccentricity up to 9 mm ($\sim 45^\circ$), though the ratio was $\sim 30\%$ greater in temporal than in nasal retina. The mRNA ratios are in good agreement with the L:M cone ratio in these same retinas, inferred from electrophysiological recordings of cone signal gain in horizontal cell interneurons. The correlation between mRNA ratios and physiological cone gain ratio supports the conclusion that both measures reflect the relative number of L and M cones. © 2000 Optical Society of America [S0740-3232(00)01403-4]

OCIS codes: 330.0330, 330.5310, 330.5380.

1. INTRODUCTION

Trichromatic color vision among primates is based on the presence of three classes of cone photoreceptor: blue [short-wavelength (S)], green [middle-wavelength (M)], and red [long-wavelength (L)] sensitive. The determination of the relative numbers and of the spatial distribution of different cone photoreceptor types in the primate retina is fundamental to our understanding of visual sensitivity and color vision. Selective staining of blue cones has allowed a direct determination of their numbers and distribution.^{1–3} While a lack of selective staining has prevented such a direct determination for L and M cones, estimates of their numbers and distribution have been derived by a variety of methods.

In primates a combined L and M cone signal forms the luminance channel in the visual pathway. The psychophysically determined human photopic luminosity function is well modeled as the weighted sum of L and M cone signals.⁴ While the mean relative L:M cone contribution to the luminosity function is $\sim 2:1$, the ratio varies greatly across individuals. This variability has been observed in a wide number of psychophysical studies.^{5–10} Estimates of the L:M cone ratio in human subjects by flicker-photometric electroretinography (ERG) have also shown wide variability in L:M ratio. In one study the L:M cone ratio varied from 0.67 to 9, with a mean of 2,¹¹ and in another study the ratio ranged from 0.28 to 8.67 with a

mean of 4:1.¹² We previously observed that the ratio of L:M mRNA present in human retinas varies widely, ranging from 1 to 10 with a mode of 4.¹³ In another study an average L:M pigment mRNA ratio of 1.5 was observed in central human retinal patches, which increased to ~ 3 for patches at 40° eccentricity.¹⁴ Microspectrophotometric measurements of eyes of individuals with normal color vision revealed an average L:M cone ratio of 1.4 in the foveal region.¹⁵ Psychophysical observations of vernier acuity were used to infer a random arrangement of L and M cones (more L than M) in the human fovea.¹⁶ Finally, the first images of the arrangement of cones in the living human eye revealed an L:M cone ratio of 1.15 in one male subject (selected for a low ratio based on ERG measurements) and an L:M cone ratio of 3.79 in a second, unselected male subject.¹⁷ The distribution of L and M cones in the subject with the 3.79 ratio was not significantly different from random distribution. The L and M cones of the subject with a ratio of 1.15 were, however, significantly more clumped than would be expected from random distribution, though a deviation from random distribution could have resulted from residual optical blur in the subject's retinal images.

In nonhuman primates estimates have generally indicated approximately equal numbers of L and M cones,^{18–20} though one exception may be the baboon retina, in which M cones appear to outnumber L cones.²¹

In the macaque monkey, ERG suggests a mean L:M cone ratio of 1:1.²² Luminance-coding ganglion cells in the macaque, however, have a spectral sensitivity that matches the human photopic luminosity function, which would suggest an L:M cone ratio of 2:1.^{23,24}

In this paper we report the L:M pigment mRNA ratios in whole retinas of three species of macaque monkey and one baboon. In several retinas we also examined the relation between the L:M pigment mRNA ratio and retinal topography by analyzing small tissue samples from known retinal locations. Finally, we compared our pigment mRNA ratio measurements with a physiological measure of the functional L:M cone ratio at the level of the horizontal cells, second-order interneurons that receive convergent input from the L and M cone population.²⁵ The results showed that, for the nonhuman primates sampled, variation in the L:M mRNA ratio was large and was comparable with that observed in the human retina. We also found a nasal-temporal asymmetry, with a reduced L:M ratio in the nasal quadrant. There was a strong correlation between the physiological and the mRNA estimates of the L:M ratio, suggesting that the two data sets are dictated by the relative numbers of L and M cones in the sample.

2. EXPERIMENTAL METHODS

A. Subjects and Retinal Specimens

Monkey retinas were obtained through the tissue distribution program of the Primate Center of the University of Washington, Seattle, Washington. The animals were unselected, did not come from the same breeding colony, and therefore were not closely related. They were free from any obvious ocular pathologies. The entire retina was dissected free of the vitreous and sclera within 1 h of death and was quickly frozen on dry ice. Punches (3 mm

in diameter) from various regions of some retinas were quickly frozen on dry ice and were kept at -70°C until they were used for RNA extraction. In some cases the retinas were kept alive for electrophysiological estimation of L and M cone ratios as described in Ref. 26 before specimens were taken for RNA analysis.

B. Extraction and Analysis of RNA

Total cellular RNA was extracted from monkey retinal tissue by the Trizol method (Gibco-BRL) and was reverse transcribed with a mixture of random hexamer primers as described in Ref. 27. First, strand cDNA was used as a template to competitively amplify, by polymerase chain reaction (PCR), segments (Fig. 1B, 215 base pairs) of the L and M pigment cDNA that encompass the 3' region of exon 4 and the 5' region of exon 5. The sequence of primers used in amplification completely matched both L and M pigment mRNA sequences of all primate species examined (Fig. 1B). Three determinations were made on three aliquots from whole retinal or from retinal patch RNA preparations.

The design of the amplification primers was based on previously determined partial coding sequences of exons 4 and 5 of L and M cone pigments of Old World monkeys.²⁸ We used this information to complete the sequences of the L and M exons 4 and 5 of *Macaca fascicularis* and *M. nemestrina* and designed primers to PCR amplify a segment encompassing the 3' region of exon 4 and the 5' region of exon 5, as shown in Fig. 1.

Primer 148 was radiolabeled at its 5' end with ^{32}P and T4 polynucleotide kinase and was used in the PCR reaction as described in Ref. 13 such that only the sense strand of the amplified products would be detectable by autoradiography after gel electrophoresis. After an initial denaturation step of 2 min at 94°C , amplification was performed for 25 cycles of denaturation at 94°C for 15 s,




Fig. 1. Strategy for amplification of the L and M exon-4-exon-5 coding region. A, Diagram of the X-chromosome-linked L-M visual pigment gene array. The numbered rectangles represent exons, and the solid lines connecting them represent introns. Diagram is not to scale. B, Sequence of the (top) M and (bottom) L segments that encompass the 3' end of exon 4 and the 5' end of exon 5, amplified with primers 148 and 78 (indicated by arrows). The dashes indicate identity in sequence between the L and the M segments. The position of intron 4 is indicated, but, because cDNA was used as the template, only exonic sequences are amplified.

followed by annealing and extension for 30 s at 64 °C. A final extension step was performed at 72 °C for 4 min. The PCR-amplified products were denatured, and the single strands corresponding to the L and M pigments (which differ in nucleotide sequence at a few positions) were resolved by electrophoresis on a nondenaturing polyacrylamide (6%) gel (20 cm × 43 cm) as described in detail elsewhere.¹³ Electrophoresis was run at a temperature of 36 °C for ~3 h. Quantification of the bands was performed by phosphorimage analysis. Because they differ in sequence (Fig. 1), the single strands of the L and the M mRNA-derived fragments assume different conformations and differ in electrophoretic mobility.

C. Correlation with Single-Cell Physiology

Horizontal cells were recorded intracellularly in an *in vitro* whole-retina preparation. The weights of L and M cone inputs to each horizontal cell were determined from the contrast gain of responses to sinusoidal flicker in different directions in color space. Detailed methods for cell identification, recording, light stimulation, data acquisition, and analysis of the L and M cone input gain to horizontal cells are given in a related paper²⁵ in which the main body of physiological results is presented. After the recordings were completed, retinal punches 3 mm in diameter centered on the location of the recorded cells were removed and were placed on dry ice until RNA extraction was started.

3. RESULTS

A. L:M Cone Pigment mRNA Ratios in Whole Retinas

First we constructed a standardization curve that related expected and observed ratios of L:M exon 4–5 sequences. Cloned and purified DNA templates that contained artificial mixtures of known ratios of purified L:M exon 4–5 sequences (labeled Input Ratios in Fig. 2) were used as templates in the PCR amplification reaction. An autoradiograph of the gel that we used to separate L from M segments and to determine their (observed) ratio is known in Fig. 2A. This assay gave a linear response of input and observed values over a wide range (0.25–8) of L:M ratios (Fig. 2B).

Next we determined the L:M mRNA ratio in whole retinas of 26 monkeys: 21 *Macaca nemestrina* (7 males, 14 females), 2 *M. fascicularis* (females), 2 *Papio anubis* (males), and a female *M. mulatta*. An autoradiograph of a single-strand conformation polymorphism gel showing separation of single strands derived from L and M exon 4–5 sequences is shown in Fig. 3A. The L:M mRNA ratios (Fig. 3B) ranged from 0.66 to 7.0, with a mean, ± 1 SD of 1.80 ± 0.60 (mean of 1.59 ± 0.56 , excluding that of monkey 6, which had a ratio of 7). The frequency distribution of the ratios is given in the inset of Fig. 3B. As was observed in human retinas,¹³ the L pigment gene is expressed at higher levels than the M pigment gene in the majority of retinas. There was no statistically significant difference in the L:M mRNA ratio between male (1.93) and female (1.77) monkeys.

B. Topographic Distribution of the L:M Cone Pigment mRNA Ratio

Next we determined the L:M cone pigment mRNA ratio in retinal punches (3 mm in diameter) distributed along the nasal–temporal and the superior–inferior axes, as shown in Fig. 4A. The results of analysis of seven eyes (six animals) are given in Table 1. The same results are shown in Fig. 4B, with the exception of that for animal 88105, for which precise eccentricity information was not available. There was no statistically significant variation in the L:M ratio with eccentricity of as much as 9 mm ($\sim 45^\circ$). On the other hand, the L:M pigment mRNA ratio in the nasal retina was consistently lower than in other regions, particularly the temporal, in 10 eyes (9 animals) (Tables 1 and 2; Fig. 4C). The nasal/temporal ratio was approximately 0.69 ± 0.14 ($p = 0.022$; two-tailed t test; $N = 10$ eyes; 9 subjects).

C. Correspondence to Physiological Measurements

In 11 instances the retinal pieces that we used to examine variation in the ratio of L:M cone pigment mRNA were derived from physiological experiments designed to measure the relative synaptic strength of L and M cone input to second-order interneurons, the H1 horizontal cells.²⁵ One finding of that study was that L:M cone input gain varies greatly as a function of retinal location and across individual retinas, similar to the L-versus-M mRNA re-




Fig. 2. Standard curve for quantification of the L:M mRNA ratio. A, Phosphor image of a single-strand conformation gel, showing the separation of L from M pigment cDNA strands. Known ratios (indicated above the lanes as input L:M mRNA ratios) of purified L and M pigment cDNA were generated and were used as templates in the competitive PCR amplification reaction by use of primers 148 (end labeled ^{32}P) and 78 (see Fig. 1B), the sequence of which matches both L and M templates. The L- and M-derived PCR products were resolved by electrophoresis, as described in Section 2. Band densities were determined with a phosphorimager. B, Plot of input and observed [± 1 standard deviation (SD)] L:M cDNA ratios.

Table 1. L:M Pigment mRNA Ratios in Monkey Retinal Sections in 3-mm Punches along the Nasal–Temporal and Superior–Inferior Axes^a

Properties	Monkey Identification Number						88105	
	89251	96111	95265	87264	91048	Left Eye	Right Eye	
	<i>M. nemestrina</i> Female	<i>M. nemestrina</i> Female	<i>M. nemestrina</i> Male	<i>M. nemestrina</i> Female	<i>M. fascicularis</i> Female	<i>M. fascicularis</i> Female	<i>M. fascicularis</i> Female	
Species	<i>M. nemestrina</i>	<i>M. nemestrina</i>	<i>M. nemestrina</i>	<i>M. nemestrina</i>	<i>M. fascicularis</i>	<i>M. fascicularis</i>	<i>M. fascicularis</i>	
Sex	Female	Female	Male	Female	Female	Female	Female	
Section								
Superior-1	2.23 ± 0.02	1.58 ± 0.26	1.13 ± 0.16	2.66 ± 0.12	1.73 ± 0.36	1.98 ± 0.22	ND	
Superior-2	1.98 ± 0.15	1.03 ± 0.21	1.33 ± 0.32	3.35 ± 0.19	1.41 ± 0.14	2.33 ± 0.25	ND	
Superior-3	2.00 ± 0.63	0.95 ± 0.26	1.78 ± 0.31	ND	2.24 ± 0.12	ND	ND	
Inferior-1	1.98 ± 0.16	0.97 ± 0.21	1.56 ± 0.25	2.41 ± 0.12	1.63 ± 0.14	2.35 ± 0.17	ND	
Inferior-2	1.95 ± 0.38	0.71 ± 0.23	1.57 ± 0.21	2.56 ± 0.06	1.39 ± 0.14	2.32 ± 0.10	ND	
Inferior-3	1.76 ± 0.02	ND	ND	ND	1.54 ± 0.72	ND	ND	
Nasal-1	1.29 ± 0.11	0.86 ± 0.04	1.57 ± 0.20	2.77 ± 0.55	1.42 ± 0.14	1.52 ± 0.11	2.26 ± 0.13	
Nasal-2	1.17 ± 0.01	0.63 ± 0.02	1.16 ± 0.12	2.21 ± 0.25	1.34 ± 0.16	1.52 ± 0.27	2.21 ± 0.19	
Nasal-3	1.22 ± 0.12	0.76 ± 0.11	1.09 ± 0.03	1.92 ± 0.10	ND	1.12 ± 0.18	3.21 ± 0.11	
Nasal-4	ND	ND	1.30 ± 0.12	ND	ND	ND	3.07 ± 0.32	
Temporal-1	2.65 ± 0.08	ND	2.15 ± 0.26	3.13 ± 0.22	1.81 ± 0.19	2.16 ± 0.10	2.90 ± 0.18	
Temporal-2	2.48 ± 0.45	ND	1.98 ± 0.23	3.39 ± 0.34	1.77 ± 0.21	2.31 ± 0.22	3.02 ± 0.17	
Temporal-3	2.56 ± 0.33	1.48 ± 0.23	ND	ND	1.61 ± 0.09	ND	3.18 ± 0.15	
Fovea	2.62 ± 0.20	1.23 ± 0.27	2.04 ± 0.34	3.13 ± 0.20	1.81 ± 0.12	1.55 ± 0.09	2.27 ± 0.14	
Remainder	2.49 ± 0.08	0.93 ± 0.25	1.35 ± 0.34	2.85 ± 0.33	1.52 ± 0.09	ND	ND	

^aEach value is the average of three independent determinations (±1 SD). ND, not determined.

Table 2. L:M Pigment mRNA Ratios in Patches from the Superior, Inferior, Temporal, and Nasal Regions^{a,b}

Sex Axis	Monkey Identification Number		
	J91076	J9026	88296
	Male	Male	Female
L-superior	1.15 ± 0.24	2.20 ± 0.24	0.86 ± 0.09
L-temporal	1.71 ± 0.04	2.38 ± 0.45	1.11 ± 0.07
L-temporal	1.85 ± 0.08	2.78 ± 0.27	ND
L-nasal	1.32 ± 0.01	1.96 ± 0.05	0.70 ± 0.06
L-nasal	ND	2.74 ± 0.16	ND
L-inferior	ND	ND	0.72 ± 0.09
Remainder	ND	2.28 ± 0.68	ND

^aEach value is the average of three independent determinations (±1 SD).

^bSpecies, *M. nemestrina*.

to determine the L:M pigment mRNA ratios in whole human retinas. This ratio ranged from 1 to 10, with a mode of 4, in Caucasian males. In the experiments reported here we applied this method to estimate the mRNA ratio in whole retinas of male and female monkeys as well as in various regions of the retina and compared these with the L:M cone ratios obtained by physiological means. As was found in human retinas,¹³ the majority of whole monkey retinas had more L than M pigment mRNA. The average L:M mRNA ratio among monkeys (the majority of monkeys were *M. nemestrina*) ranged from 0.66 to 7, with a mean of approximately 1.6 ± 0.6 . This ratio is significantly different from ratios in humans, raising the possibility that large differences in the ratio may exist among

other primate species. The mRNA ratios correlated well with the L:M cone ratios inferred from electrophysiological recordings of cone signal gain in H1 horizontal cell interneurons in the same retinas.²⁵ Thus the correlation between L:M mRNA ratios and physiological cone gain ratios supports the conclusion that both measures reflect the ratio of L:M cones in the retina. The observation that the L:M mRNA ratios in monkeys is smaller than observed among humans is consistent with previous findings by microspectrophotometry^{18,19} and ERG.²² There was more variation in the L:M ratio among the subjects in this study than was observed among the 42 male subjects (*M. mulatta* and *M. fascicularis*) studied by ERG,²² perhaps reflecting differences in the degree of genetic heterogeneity.

We observed no change in the L:M mRNA ratio in monkey retinas with eccentricities of as much as 9 mm (~45°). This result is in contrast to what was observed in 25 male human retinas, in which the L:M mRNA ratio was similar at central and mid periphery but increased substantially (by approximately a factor of 2) in far-peripheral regions (41° of eccentricity from the fovea).¹⁴ Interestingly, we observed an approximately 30% higher L:M pigment mRNA ratio in temporal than in nasal retina, indicating the presence of a temporal–nasal gradient. The significance of this gradient and how it is established would be interesting issues to explore. In contrast, no significant difference in the L:M mRNA ratio was observed between nasal and temporal patches of two male human retinas.¹⁴

It would be important to explore the mechanisms that determine the L:M cone ratio and whether the observed interindividual differences constitute a heritable trait. A mechanism that would allow the mutually exclusive expression of the X-chromosome-linked L and M pigment

genes in their respective photoreceptors has been proposed.^{13,29,30} In this mechanism a locus control region (LCR), located between 3.1 and 3.7 kbases 5' of the L pigment gene, was shown to be required for cone-photoreceptor-specific expression of both L and M pigments. Thus, if the LCR forms a stable transcriptionally active complex with the L opsin gene promoter, an L cone will result. If the LCR forms a stable complex with the proximal M pigment gene promoter, an M cone will be formed instead. Thus the choice to form an L or an M cone is predicted to be stochastic. LCR-driven mutually exclusive expression of reporter genes in transgenic mice has been demonstrated.³¹

The observed dominance of L over M cones in the majority of primate retinas may be explained in part, by the proximity of the L pigment gene to the LCR. Distance-dependent gene expression by LCR's was demonstrated for the β -globin gene locus³² and for the L-M visual pigment gene array.^{13,14,33} In addition to proximity to the LCR, differences in the sequences of the LCR, of the L and M pigment gene promoters, or of both may play a role in determining the ratio of L:M cones. This hypothesis predicts that interindividual differences in the L:M cone ratio are inherited in an X-chromosome-linked fashion. Evidence for a difference in promoter sequence that favors the M pigment gene promoter was suggested by *in vitro* promoter analysis.³⁴ Thus individual differences in the sequences of the promoter or LCR that influence the affinity of the LCR to the promoter may partly underlie individual differences in L:M cone ratio. Support for this hypothesis comes also from family studies of transmission of the L:M ratio, as estimated by photometric matches.^{6,35}




Fig. 5. Plot of percentage of L cone contrast gain for H1 horizontal cells against percentage of L pigment mRNA level for the same piece of retinal tissue. The two data sets are highly correlated (correlation coefficient $r = 0.85$; two-tailed t test; $P = 0.05$). A line with a slope of 1 is drawn through the data points. The values for the mRNA are the means of three measurements; error bars indicate ± 1 SD. Values for the H1 cell contrast gains are taken from cells located within a 3-mm-tissue punch that was used for the mRNA analysis. Multiple measurements of M and L cone contrast gain were made on each H1 cell, and the data points are mean values. SD's were small, ranging from 0.5% to 1.5% of the mean, and are not shown.

ACKNOWLEDGMENTS

The excellent technical assistance of Reiling Peng is greatly appreciated. We thank the Primate Center of the University of Washington for providing monkey retinas as part of its tissue distribution program. This research was supported by National Institutes of Health (NIH) grant EY08395 (to S. S. Deeb), by NIH grants EY06678 and EY09625 (to D. M. Dacey), and by Guggenheim Fellowship grants EY04367 and EY09625 (to D. R. Williams).

S. Deeb can be reached by e-mail at deeb@genetics.washington.edu.

REFERENCES

1. F. M. de Monasterio, E. P. McCrane, J. K. Newlander, and S. J. Schein, "Density profile of blue-sensitive cones along the horizontal meridian of macaque retina," *Invest. Ophthalmol. Visual Sci.* **26**, 289–302 (1985).
2. C. A. Curcio, K. A. Allen, K. R. Sloan, C. L. Lerea, J. B. Hurley, I. B. Klock, and A. H. Milam, "Distribution and morphology of human cone photoreceptors stained with anti-blue opsin," *J. Comp. Neurol.* **312**, 610–624 (1991).
3. A. Szel, T. Diamantstein, and P. Rohlich, "Identification of the blue-sensitive cones in the mammalian retina by anti-visual pigment antibody," *J. Comp. Neurol.* **273**, 593–602 (1988).
4. V. C. Smith and J. Pokorny, "Spectral sensitivity of the foveal cone photopigments between 400 and 500 nm," *Vision Res.* **15**, 161–171 (1975).
5. W. A. Rushton and H. D. Baker, "Red/green sensitivity in normal vision," *Vision Res.* **4**, 75–85 (1964).
6. H. L. De Vries, "The heredity of the relative numbers of red and green receptors in the human eye," *Genetica (The Hague)* **24**, 199–212 (1948).
7. R. L. Vimal, J. Pokorny, V. C. Smith, and S. K. Shevell, "Foveal cone thresholds," *Vision Res.* **29**, 61–78 (1989).
8. M. F. Wesner, J. Pokorny, S. K. Shevell, and V. C. Smith, "Foveal cone detection statistics in color-normals and dichromats," *Vision Res.* **31**, 1021–1037 (1991).
9. C. M. Cicerone and J. L. Nerger, "The relative numbers of long-wavelength-sensitive to middle-wavelength-sensitive cones in the human fovea centralis," *Vision Res.* **29**, 115–128 (1989).
10. J. L. Nerger and C. M. Cicerone, "The ratio of L cones to M cones in the human parafoveal retina," *Vision Res.* **32**, 879–888 (1992).
11. J. Jacobs and J. Neitz, "Electrophysiological estimates of individual variation in the L/M cone ratio," in *Proceedings of the International Symposium on Color Vision Deficiencies*, B. Drum, ed. (Kluwer Academic, Dordrecht, The Netherlands, 1993), Vol. VI, pp. 107–112.
12. J. Kremers, T. Usui, H. P. Scholl, and L. T. Sharpe, "Cone signal contributions to electrograms in dichromats and trichromats," *Invest. Ophthalmol. Visual Sci.* **40**, 920–930 (1999).
13. T. Yamaguchi, A. G. Motulsky, and S. S. Deeb, "Visual pigment gene structure and expression in human retinae," *Hum. Mol. Genet.* **6**, 981–990 (1997).
14. S. A. Hagstrom, J. Neitz, and M. Neitz, "Variations in cone populations for red-green color vision examined by analysis of mRNA," *NeuroReport* **9**, 1963–1967 (1998).
15. H. J. A. Dartnall, J. K. Bowmaker, and J. D. Mollon, "Human visual pigments: microspectrophotometric results from the eyes of seven persons," *Proc. R. Soc. London* **220**, 115–130 (1983).
16. P. D. Gowdy and C. M. Cicerone, "The spatial arrangement of the L and M cones in the central fovea of the living human eye," *Vision Res.* **38**, 2575–2589 (1998).
17. A. Roorda and D. R. Williams, "The arrangement of the

- three cone classes in the living human eye," *Nature* **397**, 520–522 (1999).
18. J. K. Bowmaker, "Microspectrophotometry of vertebrate photoreceptors. A brief review," *Vision Res.* **24**, 1641–1650 (1984).
 19. J. D. Mollon and J. K. Bowmaker, "The spatial arrangement of cones in the primate fovea," *Nature* **360**, 677–679 (1992).
 20. O. S. Packer, D. R. Williams, and D. G. Bensinger, "Photopigment transmittance imaging of the primate photoreceptor mosaic," *J. Neurosci.* **16**, 2251–2260 (1996).
 21. R. E. Marc and H. G. Sperling, "Chromatic organization of primate cones," *Science* **196**, 454–456 (1977).
 22. G. H. Jacobs and J. F. Deegan II, "Spectral sensitivity of macaque monkeys measured with ERG flicker photometry," *Visual Neurosci.* **14**, 921–928 (1997).
 23. B. B. Lee, P. R. Martin, and A. Valberg, "The physiological basis of heterochromatic flicker photometry demonstrated in the ganglion cells of the macaque retina," *J. Physiol. (London)* **404**, 323–347 (1988).
 24. R. Shapley, "Parallel neural pathways and visual function," in *The Cognitive Neurosciences*, M. S. Gazzaniga, E. Bizzi, I. B. Black, C. Blakemore, L. Cosmides, S. M. Kosslyn, J. E. LeDoux, J. A. Movshon, S. Pinker, M. I. Posner, P. Rakic, D. L. Schacter, J. Tooby, and E. Tulving, eds. (MIT Press, Cambridge, Mass., 1995), pp. 315–324.
 25. D. M. Dacey, L. C. Diller, J. Verweij, and D. R. Williams, "Physiology of L- and M-cone inputs to H1 horizontal cells in the primate retina," *J. Opt. Soc. Am. A* **17**, 589–596 (2000).
 26. D. M. Dacey, "Circuitry for color coding in the primate retina," *Proc. Natl. Acad. Sci. USA* **93**, 582–588 (1996).
 27. J. Winderickx, L. Battisti, Y. Hibiya, A. G. Motulsky, and S. S. Deeb, "Haplotype diversity in the human red and green opsin genes: evidence for frequent sequence exchange in exon 3," *Hum. Mol. Genet.* **2**, 1413–1421 (1993).
 28. R. E. Ibbotson, D. M. Hunt, J. K. Bowmaker, and J. D. Mollon, "Sequence divergence and copy number of the middle- and long-wave photopigment genes in Old World monkeys," *Proc. R. Soc. London Ser. B* **247**, 145–154 (1992).
 29. Y. Wang, J. P. Macke, S. L. Merbs, D. J. Zack, B. Klaunberg, J. Bennett, J. Gearhart, and J. Nathans, "A locus control region adjacent to the human red and green visual pigment genes," *Neuron* **9**, 429–440 (1992).
 30. J. Winderickx, L. Battisti, A. G. Motulsky, and S. S. Deeb, "Selective expression of human X chromosome-linked green opsin genes," *Proc. Natl. Acad. Sci. USA* **89**, 9710–9714 (1992).
 31. Y. Wang, P. M. Smallwood, M. Cowan, D. Blesh, A. Lawler, and J. Nathans, "Mutually exclusive expression of human red and green visual pigment reporter transgenes occurs at high frequency in murine cone photoreceptors," *Proc. Natl. Acad. Sci. USA* **96**, 5251–5256 (1999).
 32. F. Grosveld, E. de Boer, N. Dillon, J. Gribnau, T. McMorrow, E. Milot, T. Trimborn, M. Wijgerde, and P. Fraser, "The dynamics of globin gene expression and position effects," *Novartis Found. Symp.* **214**, 67–79 (1998).
 33. T. Hayashi, A. G. Motulsky, and S. S. Deeb, "Position of a 'green-red' hybrid gene in the visual pigment array determines colour-vision phenotype," *Nat. Genet.* **22**, 90–93 (1999).
 34. S. A. Shaaban and S. S. Deeb, "Functional analysis of the promoters of the human red and green visual pigment genes," *Invest. Ophthalmol. Visual Sci.* **39**, 885–896 (1998).
 35. M. Lutze, N. J. Cox, V. C. Smith, and J. Pokorny, "Genetic studies of variation in Rayleigh and photometric matches in normal trichromats," *Vision Res.* **30**, 149–162 (1990).